

## Emergency Contraception

Adapted from Women and HIV: A Case Based Focus on Contraception, Family Planning, Prenatal Care and Postpartum Care - FL APPROVED -

By Melissa Badowski, PharmD

Regimen	Available Formulation	Timing of Use After Unprotected Encounter	Access	Interactions with ARVs
Selective progesterone receptor modulator	<ul style="list-style-type: none"> <li>• Ulipristal acetate 30 mg</li> <li>• One tablet</li> </ul>	Up to 5 days	Prescription	Contraindicated Cobicistat, Efavirenz, and PIs
Progestin only	<ul style="list-style-type: none"> <li>• Levonorgestrel 1.5 mg</li> <li>• One tablet</li> </ul>	Up to 3 days	OTC without age restriction	Avoid Cobicistat, Efavirenz, and PIs
	<ul style="list-style-type: none"> <li>• Levonorgestrel 0.75 mg</li> <li>• Two tablets</li> </ul>		OTC for those $\geq 17$ years with photo ID	
Combined progestin-estrogen pills	<ul style="list-style-type: none"> <li>• Various formulations</li> <li>• <a href="http://ec.princeton.edu/questions/dose.html#dose">http://ec.princeton.edu/questions/dose.html#dose</a></li> </ul>	Up to 5 days	Prescription	Avoid Cobicistat, Efavirenz, and PIs
Copper Intrauterine Device (IUD)	<ul style="list-style-type: none"> <li>• IUD</li> </ul>	Up to 5 days	Office visit with clinician	None

Resources:

- ACOG. 2015;126(3):e1-11.