

## Naloxone Dosing

Adapted from Managing Opioid Overdose in the Community

By Bethany DiPaula, PharmD, BCPP

Formulation	Instructions for Use
Naloxone Intranasal • 2mg/2ml	<b>For intranasal administration</b> 2mg/2mL single-dose Luer-Jet prefilled syringe. Include one luer-lock mucosal atomization device per dose dispensed. <b>Directions for use:</b> Spray 1 mL in each nostril. Repeat after 3 minutes if no or minimal response
Naloxone Intranasal (Narcan®) • 2mg/0.4ml • 4mg/0.4ml	<b>For intranasal administration</b> NARCAN® 4 mg/0.4mL nasal spray. Include <b>Directions for use:</b> Administer a single spray of NARCAN® in one nostril. Repeat after 3 minutes if no or minimal response.
Naloxone Intramuscular • 0.4mg/ml	<b>For intramuscular injection</b> 0.4mg/mL in 1mL single dose vials. Include one 3cc, 23g, 1" syringe per dose dispensed. <b>Directions for use:</b> Inject 1 mL IM in shoulder or thigh. Repeat after 3 minutes if no or minimal response.
Naloxone (Evzio®) • 0.4mg/0.4ml (d/c) • 2mg/0.4ml	<b>For intramuscular or subcutaneous injection</b> EVZIO® 2mg/0.4mL auto-injector, #1 Two-pack <b>Directions for use:</b> Follow audio instructions from device. Place on thigh and inject 0.4 mL. Repeat after 3 minutes if no or minimal response.

Reference(s):

- [http://prescribetoprevent.org/wp2015/wp-content/uploads/Naloxone-product-chart.16\\_01\\_21.pdf](http://prescribetoprevent.org/wp2015/wp-content/uploads/Naloxone-product-chart.16_01_21.pdf)